

BONDAGE NUMBER FOR JUMP GRAPH OF SOME TREES

Betül Atay

Communicated by Mehmet Gümüş

Abstract: The dominating set of a graph is a vertex set in that every vertex which is not in the dominating set is adjacent to at least one vertex of the dominating set. The domination number is the minimal cardinality among all dominating sets. The bondage number of any graph is the minimal cardinality among all sets of edges whose removal from the graph results in a graph with domination number greater than the domination number of the preliminary graph. In this paper, we investigate the bondage number for jump graph of some certain trees.

Keywords and phrases: Domination number, bondage number, jump graph.

MSC 2010 Classifications: Primary 05C05; Secondary 05C69.

1 Introduction

Graph theory has become one of the most powerful mathematical tools in the analysis and study of the architecture of networks whose vertices represent the components of the system and the edges represent connection between a pair of vertices that enable mutual communication. The vulnerability of a communication network measures the resistance of network to the disruption of operation after the failure of certain stations or communication links. For any communication network greater degrees of stability or less vulnerability is required. Vulnerability can be measured by certain parameters like domination, bondage number, connectivity, betweenness, binding number, toughness, scattering number, integrity etc. Graph theory is among the popular methods for solving many complex problems. Graph theory increases its development and usage area due to the easy modelling of daily problems and successful results of effective solution methods. The dominant nodes indicate the dominance of the people or objects modelled on the graph over each other. However, minimum domination set aims to connect all vertices in the graph with the least number of vertices selected on the graph. Determining the minimum dominating set in graphs is one of the most difficult problems defined as NP-hard. When the usage areas of dominating sets in graphs examined, it is seen to provide significant gains in many areas such as social networks, transportation systems, telecommunication, defence industry, health systems, etc.

In a graph $G = (V(G), E(G))$, a subset $S \subseteq V(G)$ of vertices is a *dominating set* if every vertex in $V(G) - S$ is adjacent to at least one vertex of S . The *domination number* $\gamma(G)$ is the minimal cardinality of a dominating set. One of the vulnerability parameters based on domination number known as bondage number in a graph G examines the situation in which the domination number increases if some connections are broken. The opposite parameter, that is, examining the decrease in the domination number, is the reinforcement number. The *bondage number* $b(G)$ of a nonempty graph G is the cardinality of a smallest set of edges whose removal from G results in a graph with domination number greater than $\gamma(G)$. That is,

$$b(G) = \min\{|S| : S \subseteq E(G), \gamma(G - S) > \gamma(G)\}.$$

We call such an edge set S that $\gamma(G - S) > \gamma(G)$ the *bondage set* and the minimum one the *minimum bondage set*. If $b(G)$ does not exist, for example empty graphs, then $b(G) = \infty$ is defined.

The bondage number was introduced by Bauer, Harary, Nieminen and Suffel [3], and has been further studied by Fink, Jacobson, Kinch and Roberts[8], Hartnell and Rall[4] and others. Later, the bondage number for middle graphs and complementary prism graphs was studied in [1, 2].

In this paper, the graph G is taken as a simple, finite and undirected graph with vertex set $V(G)$ and edge set $E(G)$. The distance $d(u, v)$ between two vertices u and v in G is the length of a shortest path joining them if any; otherwise $d(u, v) = \infty$. A shortest $u-v$ path is often called a *geodesic*. The *diameter* of G , denoted by $diam(G)$ is the largest distance between two vertices in $V(G)$. The number of the neighbor vertices of the vertex v is called degree of v and denoted by $deg_G(v)$. The minimum and maximum degrees of a vertex of G are denoted by $\delta(G)$ and $\Delta(G)$. A vertex v is said to be pendant vertex if $deg_G(v) = 1$. A vertex u is called support if u is adjacent to a pendant vertex [5]. Let u be a vertex of a graph $G = (V, E)$. Then $N(u) = \{v \in V(G), v \text{ and } u \text{ are adjacent}\}$ is the open neighborhood of u , and $N[u] = \{u\} \cup N(u)$ denotes the closed neighborhood of u . The *eccentricity* $e(v)$ of a vertex v in a connected graph G is $\max d(u, v)$ for all u in G . The *radius* $r(G)$ is the minimum eccentricity of the vertices. Note that the maximum eccentricity is the diameter. A vertex v is a central vertex if $e(v) = r(G)$, and the *center* of G is the set of all central vertices [5]. The connectivity $\kappa = \kappa(G)$ of a graph G is the minimum number of vertices whose removal results in a disconnected or trivial graph. A vertex and an edge are said to *cover* each other if they are incident. A set of vertices which covers all the edges of a graph G is called a *vertex cover* for G , while a set of edges which covers all the vertices is a *edge cover*. The smallest number of vertices in any vertex cover for G is called its *vertex covering number* and is denoted by $\alpha_0(G)$ or α_0 . Similarly, $\alpha_1(G)$ or α_1 is the smallest number of edges in any edge cover of G and is called its *edge covering number*. A set of vertices in G is independent if no two of them are adjacent. The largest number of vertices in such a set is called the *vertex independence number* of G and is denoted by $\beta_0(G)$ or β_0 . Analogously, an independent set of edges of G has no two of its edges adjacent and the maximum cardinality of such a set is the *edge independence number* $\beta_1(G)$ or β_1 [5]. The complement \overline{G} of a simple graph G is the simple graph with vertex set $V(G)$ defined by $uv \in E(\overline{G})$ if and only if $uv \notin E(G)$. The corona $(G_1 \circ G_2)$ of two graphs G_1 and G_2 is defined as the graph G obtained by taking one copy of G_1 (which has p_1 vertices) and p_1 copies of G_2 , and then joining the i th vertex of G_1 to every vertex in the i th copy of G_2 [11]. Let G_1 and G_2 be two graphs with vertex sets are $V(G_1)$ and $V(G_2)$; edge sets are $E(G_1)$ and $E(G_2)$ respectively. Then, the join of $G_1 + G_2$ of two graphs G_1 and G_2 is the graph with vertex set $V(G_1 + G_2) = V(G_1) \cup V(G_2)$ and edge set $E(G_1 + G_2) = E(G_1) \cup E(G_2) \cup \{uv : u \in V(G_1), v \in V(G_2)\}$ [13]. The line graph $L(G)$ of G has the edges of G as its vertices which are adjacent in $L(G)$ if and only if the corresponding edges are adjacent in G . We call the complement of line graph $L(G)$ as the jump graph $J(G)$ of G . The jump graph $J(G)$ of a graph G is the graph defined on $E(G)$ and in which two vertices are adjacent if and only if they are not adjacent in G . Since both $L(G)$ and $J(G)$ are defined on the edge set of a graph G , it follows that isolated vertices of G (if G has) play no role in line graph and jump graph transformation. We assume that the graph G under consideration is nonempty and has no isolated vertices [6, 14]. Aim of the paper is to establish the bondage number for jump graph of some certain trees. Graph operations like join, corona, line and jump are play important role in obtaining larger structures from small structures. Moreover, trees are more consistent than general graph structures since each edge is an intersection edge. Undirected trees are the simplest yet most important class of graphs for computing domination and bondage numbers. Therefore, the study of bondage number for jump graphs of trees may be useful in studying other graph families.

2 Basic Results

In this section, some well-known basic results are given with regard to domination and bondage number.

Theorem 2.1. [8] For a complete graph K_n of order $n \geq 2$, then $b(K_n) = \lceil \frac{n}{2} \rceil$.

Theorem 2.2. [8] For a path graph P_n of order $n \geq 2$, then

$$b(P_n) = \begin{cases} 2, & \text{if } n \equiv 1 \pmod{3} \\ 1, & \text{otherwise} \end{cases}$$

Theorem 2.3. [8] For a cycle graph C_n of order $n \geq 3$, then

$$b(C_n) = \begin{cases} 3, & \text{if } n \equiv 1 \pmod{3} \\ 2, & \text{otherwise} \end{cases}$$

Theorem 2.4. [8] For a star graph $K_{1,n}$ of order $n + 1$, where $n \geq 2$. Then, $b(K_{1,n}) = 1$.

Theorem 2.5. [8] If G is a nonempty graph with a unique minimum dominating set, then $b(G) = 1$.

Theorem 2.6. [8] If T is a nontrivial tree, then $b(T) \leq 2$.

Theorem 2.7. [8] If G is a connected graph of order $n \geq 2$, then $b(G) \leq n - 1$.

Theorem 2.8. [8] If G is a nonempty graph, then

$$b(G) \leq \min\{\deg(u) + \deg(v) - 1 : u \text{ and } v \text{ are adjacent vertices.}\}$$

Theorem 2.9. [8] If $\Delta(G)$ and $\delta(G)$ denote respectively the maximum and minimum degree among all vertices of nonempty connected graph G , then $b(G) \leq \Delta(G) + \delta(G) - 1$.

Theorem 2.10. [8] If G is a nonempty graph with domination number $\gamma(G) \geq 2$, then $b(G) \leq (\gamma(G) - 1)\Delta(G) + 1$.

Theorem 2.11. [8] If G is a connected graph of order $n \geq 2$, then $b(G) \leq n - \gamma(G) + 1$.

Theorem 2.12. [8] If G is a nonempty graph, then $b(G) \leq \Delta(G) + 1$.

Theorem 2.13. [4] If G has edge connectivity κ , then $b(G) \leq \Delta(G) + \kappa - 1$.

Theorem 2.14. [2] If G is a nonempty connected graph of order n with $\alpha_0(G) = 1$, then $b(G) = 1$.

Theorem 2.15. [14] Let $G(p, q)$ be a graph with $p = |V|$ and $q = |E|$, then

- a) For any connected (p, q) graph G , $\gamma(J(G)) \leq q - \beta_1(G) + 1$.
- b) For any connected (p, q) graph G , $\gamma(G) + \gamma(J(G)) < (\frac{p+1}{2})^2$.
- c) For any connected graph G with diameter, $\text{diam}(G) \geq 2$, $\gamma(J(G)) \geq 2$.
- d) For any tree T with diameter greater than 3, $\gamma(J(T)) = 2$.
- e) For any connected (p, q) graph G , $\gamma(J(G)) \geq q - \Delta(G)$
- f) For any connected (p, q) graph G , $2 \leq \gamma(J(G)) \leq \lfloor \frac{q}{2} \rfloor$.
- g) For any connected graph G without pendant vertex, $\gamma(J(G)) \leq \delta(G)$.

3 Bondage Number for Jump Graph of Some Trees

In this section, we calculated the bondage number for jump graph of some trees as comet graph $C_{n,t}$, k -ary H_n^k , caterpillar $T_{n,m}$, the graph E_n^t and banana tree $B_{n,k}$.

Definition 3.1. [9] For integer $n \geq 2$ and $t \geq 1$, the comet graph $C_{n,t}$ is defined to be the graph of order $n + t + 1$ obtained from disjoint union of a star $K_{1,n}$ and a path P_t with t vertices by adding an edge joining the central vertex of the star with an end-vertex of the path. A Comet graph $C_{5,3}$ is illustrated in Figure 1.




Figure 1. Comet graph $C_{5,3}$

Theorem 3.2. *The bondage number for the jump graph of comet graph $C_{n,t}$ is $b(J(C_{n,t})) = t - 2$.*

Proof. We know from the Theorem 2.15(d) that $\gamma(J(C_{n,t})) = 2$. We can label the corresponding vertices to the relevant edges in the comet graph $C_{n,t}$ as $V(J(C_{n,t})) = \{c_1, c_2, \dots, c_n, c_{n+1}, (n+1)(n+2), (n+2)(n+3), \dots, (t-2)(t-1), (t-1)t\}$. There are n vertices c_1, c_2, \dots, c_n degree of $t-1$, one vertex c_{n+1} degree of $t-2$, $t-2$ vertices $(n+1)(n+2), (n+2)(n+3), \dots, (t-2)(t-1)$ degree of $n+t-3$ and one vertex $(t-1)t$ degree of $n+t-2$ in $J(C_{n,t})$. If we delete the edges that are incident to the vertex c_{n+1} with minimum degree, then the graph formed by join of the complement graph $\overline{P_{t-1}}$ and n disjoint vertices means the complement graph $\overline{K_n}$ and one isolated vertex c_{n+1} remain. If we denote the set of the deleted edges by S , then $\gamma(J(C_{n,t}) - S) = \gamma(\overline{K_n} + \overline{P_{t-1}}) = 2$, except the vertex c_{n+1} . Since, c_{n+1} dominates itself, the domination number of the remaining graph results with 3, which is one more than $\gamma(J(C_{n,t}))$. Hence, we have $b(J(C_{n,t})) = t - 2$ because of $deg(c_{n+1}) = t - 2$. This completes the proof. \square

Definition 3.3. [10] The complete $k - ary$ tree H_n^k of depth n is the rooted tree in which all vertices at level $n - 1$ or less have exactly k children, and all vertices at level n are leaves. A complete $2 - ary$ tree H_4^2 is illustrated in Figure 2.




Figure 2. 2-ary tree H_4^2

Theorem 3.4. *The bondage number for the jump graph of $k - ary$ tree H_n^k is $b(J(H_n^k)) = \frac{k^n - k}{k-1} \binom{k^n + k - 2}{k-1}$.*

Proof. The Line graph of H_n^k is the graph formed by combining the root vertices of two graphs H_{n-1}^k . The jump graph $J(H_n^k)$, which is the complement of line graph $L(H_n^k)$ is a graph whose root vertex of one of H_{n-1}^k is adjacent to all vertices except the root vertex of the other graph H_{n-1}^k . All other vertices of these two components are adjacent to each other. That is, the root vertex of H_{n-1}^k is adjacent to the all remaining vertices $A = k + k^2 + k^3 + \dots + k^{n-1} = \frac{k^n - k}{k-1}$ of H_{n-1}^k ; are adjacent to $A + 1 = \frac{k^n - 1}{k-1}$ vertices including the root vertex of the other component H_{n-1}^k . We know $\gamma(J(H_{n-1}^k)) = 2$ from the Theorem 2.15(d). Let S be the set of edges connecting these two components. So, if we disconnect these two components, which are the same, then $\gamma(J(H_n^k) - S) = 2\gamma(J(H_{n-1}^k)) = 2 + 2 = 4 > 2 = \gamma(J(H_n^k))$. In order to achieve this, edges equal to the cardinality of the edge set S must be removed from the $J(H_n^k)$. Hence, $b(J(H_n^k)) = A^2 + 2A = \left(\frac{k^n - k}{k-1}\right)^2 + 2\frac{k^n - k}{k-1} = \frac{k^n - k}{k-1} \binom{k^n + k - 2}{k-1}$ is obtained and this completes the proof. \square

Definition 3.5. [7] A tree T is called a caterpillar, if removal of all its pendant vertices results in a path called the spine of T , denoted by $sp(T)$. If all vertices of $sp(T)$ have equal number of pendant vertices, then the resulting graph is called a regular caterpillar. A regular caterpillar can also be defined as the corona of two special graph types. That is, if $T_{n,m}$ is a regular caterpillar of order $nm + n$, then $T_{n,m} \cong P_n \circ mK_1$. A regular caterpillar $T_{5,3}$ is illustrated in Figure 3.




Figure 3. Caterpillar graph $T_{5,3}$

Theorem 3.6. *The bondage number for the jump graph of regular caterpillar graph $T_{n,m}$ is $b(J(T_{n,m})) = nm + n - 2m - 4$.*

Proof. Spines of line graph and its complement jump graph of regular caterpillar $T_{n,m}$ contains $n - 1$ vertices. The vertex which has maximum degree in $L(T_{n,m})$ is the vertex with minimum degree in $J(T_{n,m})$. If we delete the edges that are incident to the vertex with minimum degree in $J(T_{n,m})$, then two graphs $J(T_{t,m})$ and $J(T_{n-t,m})$ remain. We know $\gamma(J(T_{t,m})) = \gamma(J(T_{n-t,m})) = 2$. As the isolated vertex denoted by v dominates itself, the domination number of the remaining graph results with 5, which is 3 more than $\gamma(J(T_{n,m}))$. Since, $deg(v) = 2t + 2$ in $L(T_{n,m})$ and $deg(v) = nm + n - 2m - 4$ in $J(T_{n,m})$, $b(J(T_{n,m})) = nm + n - 2m - 4$ is obtained. This completes the proof. \square

Definition 3.7. [12] The graph E_n^t is a tree which has t legs and each leg has n vertices. Thus, E_n^t has $nt + 2$ vertices. The graph E_4^3 is illustrated in Figure 4.




Figure 4. The graph E_4^3

Theorem 3.8. *The bondage number for the jump graph of E_n^t is $b(J(E_n^t)) = (n - 1)t$.*

Proof. $L(E_n^t)$ is the graph formed by combining the one of the pendant vertices of each path graph P_n with the vertex uv . Hence, there are the vertex uv degree of t , $(n - 1)t$ vertices degree of 2 and t pendant vertices in $L(E_n^t)$. The vertex with the minimum degree is the vertex uv in $J(E_n^t)$ and $deg(uv) = (n - 1)t$. By definition, the γ -set includes the vertices of maximum degree. In order to increase the cardinality of the γ -set, the edge set S , which consists the edges that are incident to the vertex uv must be remove from $J(E_n^t)$. Therefore, $\gamma(J(E_n^t) - uv) = 2$ and uv must dominate itself. Hence, $\gamma(J(E_n^t) - S) = 3$, which is 1 more than $\gamma(J(E_n^t))$. This completes the proof. \square

Definition 3.9. [15] Banana tree is obtained by linking one leaf of each of n copies of an k vertices star graph structure with a single root vertex that is distinct from all the stars and the tree is denoted by $B_{n,k}$. The Banana tree $B_{3,5}$ is illustrated in Figure 5.




Figure 5. Banana tree $B_{3,5}$

Theorem 3.10. *The bondage number for the jump graph of banana tree $B_{n,k}$ is*

$$b(J(B_{n,k})) = \begin{cases} nt - t, & \text{if } n < t \\ nt - n - 1, & \text{if } n \geq t \end{cases}$$

Proof. As in the previous proof, when the minimum degree vertex in $J(B_{n,k})$ is isolated, the bondage number of the remaining graph is 2. Let's denote the vertex with the minimum degree by v . If $n < t$, then $\deg(v) = nt - t$ and if $n \geq t$, then $\deg(v) = nt - n - 1$. Hence, we have

$$b(J(B_{n,k})) = \begin{cases} nt - t, & \text{if } n < t \\ nt - n - 1, & \text{if } n \geq t \end{cases}$$

This completes the proof. □

4 Conclusion

In this paper, we studied the bondage number for jump graphs of some trees including Comet graph $C_{n,t}$, k -ary tree H_n^k , caterpillar graph $T_{n,m}$, the graph E_n^t and banana tree $B_{n,k}$. General results were obtained for these structures. In the graphs examined, it was observed that the bondage number was obtained by splitting the graph into two components or by leaving an isolated vertex. As one of the most relevant application areas, a problem related to social networks was presented. Thus, the increase in the robustness and stability of the given graphs by breaking some links was investigated.

Acknowledgment

The author expresses her sincere gratitude to the area editor and reviewer for useful advice.

References

- [1] A.Aytaç and T.Turacı, On the bondage, strong and weak bondage numbers in complementary prism graphs, *Computer Science Journal of Moldova*, **29:1(85)**, 59–75 (2021).
- [2] A. Aytaç, T.Turacı and Z.N. Odabaş, On the bondage number of middle graphs, *Mathematical Notes*, **93:6**, 795–801 (2013).
- [3] D. Bauer, F.Harary, J. Nieminen and C.L. Suffel, Domination alteration sets in graph, *Discrete Math*, **47:2-3**, 153–161 (1983).
- [4] D.F.Hartnell and D.F.Rall, Bounds on the bondage number of a graph, *Discrete Math*, **128:1-3**, 173–177 (1994).
- [5] F. Harary, Graph Theory, *Addition-Wesley Publishing Co., Reading, MA/Menlo Park, CA/London*, (1969).
- [6] G. Chartrand, H. Hevia, E.B. Jarrett and M. Schultz, Subgraph distances in graphs defined by edge transfers, *Discrete Math.*, **170**, 63–79 (1997).
- [7] J.A. Gallian, A dynamic survey of graph labelling, *Elect. Jour. Combin.*, 15, DS6. (2008).
- [8] J.F.Fink, M.S.Jacobson, L.F. Kinch and J.Roberts, The bondage number of a graph, *Discrete Math*, **86:1-3**, 47-57 (1990).
- [9] M. Symen, M. Plipczuk and R. Skrekovski, Relation between Randic index and average distance of trees, *Institute of Mathematics, Physics and Mechanics Jadranska*, 48, Slovenia, 1130 (2010).
- [10] P. Dankelmann, D. Day, D. Erwin, S. Mukwembi and H. Swart, Domination with exponential decay, *Discrete Mathematics*, 309, 5877–5883 (2009).
- [11] R. Frucht and F. Harary, On the corona two graphs, *Aequationes Math.*, **4**, 322–325 (1970).
- [12] T. Cormen, C.E. Leiserson and R.L. Rivest, *Introduction to Algorithms*, the MIT Pres. (Fourth edition), (1990).
- [13] T.W. Haynes, S.T. Hedetniemi and P.J. Slater, *Fundamentals of domination in graphs*, *Markel Dekker, Inc., New York*
- [14] Y.B. Maralabhavi, S.B. Anupama and M. Goudar Venkanagouda, Domination number of jump graph, *International Mathematical Forum*, **8:16**, 753–758 (2013).

- [15] W.-C. Chen, H.-I. Lu and Y.-N. Yeh, Operations of Interlaced Trees and Graceful Trees, *Southeast Asian Bull. Math.* 21, 337–348 (1997).

Author information

Betül Atay, Department of Computer Education and Instructional Technology, Ağrı İbrahim Çeçen University, 04100, Ağrı, TURKEY.

E-mail: batay@agri.edu.tr

Received: 25.04.2024

Accepted: 31.05.2024

Published: 30.06.2024